

Observation of Coherent Exclusive ρ^0 Production in Ultra-Peripheral Relativistic Heavy Ion Collisions

Falk Meissner, Spencer Klein, Vladimir Morozov,
Akio Ogawa*, Janet Seger†, Jim Thomas, Pablo Yepes‡ and the STAR Collaboration

We present the first observation of coherent ρ^0 production in ultra-peripheral relativistic heavy ion collisions. At impact parameters $b > 2R_A$ the nuclei do not physically collide, but interact via long-range electromagnetic fields. Two photo-nuclear processes are observed at STAR: exclusive ρ^0 production, $AuAu \rightarrow AuAu\rho^0$, and ρ^0 production with accompanying nuclear excitation, $AuAu \rightarrow Au^*Au^*\rho^0$. To study the later process, a data set of about 800k 'minimum bias' events at $\sqrt{S_{NN}} = 130$ GeV/nucleon is used, where the coincident detection of neutrons from nuclear break up in the zero degree calorimeters is required as a trigger.

Events were selected with exactly two tracks that formed a primary vertex. The pair transverse momentum spectrum in Fig. 1a) is peaked at $p_T \sim 50$ MeV, which is the signature for coherent coupling to both nuclei.¹ A normalized background model from like-sign combination pairs (histogram) does not show such a peak.

The acceptance and efficiency corrected $M_{\pi\pi}$ invariant mass spectrum of Fig. 1b) shows a clear peak at the ρ^0 mass for the coherent events with a pair $p_T < 150$ MeV. The shaded histogram is the combinatorial background; the hashed histogram contains additionally the contribution from coherent e^+e^- pairs.² The spectrum is fitted (solid line) by a relativistic Breit-Wigner for the ρ , a mass-independent contribution from direct $\pi^+\pi^-$ production (both dashed), and their interference.³ A polynomial contribution for the

background (dash-dotted) is included in the fit. The ρ^0 mass and width are consistent with the known values. The amplitude ratio of direct pion pair to ρ production is comparable to photon-proton scattering data. Preliminary, the cross section for coherent ρ^0 production accompanied by nuclear excitation $\sigma(AuAu \rightarrow Au_{xn}^*Au_{xn}^*\rho^0)$ is of order 30 mb and the cross section for ρ^0 production with nuclear breakup under emission of single neutrons $\sigma(AuAu \rightarrow Au_{1n}^*Au_{1n}^*\rho^0)$ is of order 2 mb. Here, the nuclear excitation is dominated by the Giant Dipole Resonance.




Figure 1: (a) The p_T distribution for ρ candidates from the minimum bias data sample. (b) The $M_{\pi\pi}$ invariant mass distribution for $p_T < 0.15$ GeV/c.

*Penn. State. Univ, University Park, PA.

†Creighton Univ., Omaha, NE.

‡Rice University, Houston, TX.

¹S. Klein, J. Nystrand, Phys. Rev. **C60**, 014903 (1999).

²F. Meissner, in *Proc. Photon 2001*, nucl-ex/0112008, F. Marozov *et. al*, this volume.

³ P. Söding, Phys. Lett. **19** 702 (1966), and Phys. Rev.

Lett. **22** 981 (1969), T.H. Bauer *et al.*, Rev. Mod. Phys. **50** 261 (1978).